

On embedding all n -manifolds into a single $(n + 1)$ -manifold

Fan Ding, Shicheng Wang and Jiangang Yao

February 2, 2008

Abstract

For each composite number $n \neq 2^k$, there does not exist a single connected closed $(n + 1)$ -manifold such that any smooth, simply-connected, closed n -manifold can be topologically flat embedded into it. There is a single connected closed 5-manifold W such that any simply-connected, 4-manifold M can be topologically flat embedded into W if M is either closed and indefinite, or compact and with non-empty boundary.

1 Introduction and some prerequisites

The celebrated Whitney embedding theorem states that any smooth n -dimensional manifold can be embedded smoothly into the Euclidean space $\mathbb{R}^{2n}$, or equivalently, any smooth n -dimensional manifold can be embedded into the sphere S^{2n} , the simplest closed $2n$ -manifold. If the target space is allowed to be other closed manifolds, it is natural to wonder the following general problem:

Find the smallest nonnegative integer e_n , such that any n -dimensional connected, closed manifold can be (topologically flat or smoothly) embedded into a single connected, closed manifold of dimension $n + e_n$.

Clearly $0 \leq e_n \leq n$.

Examples. The only known e_i 's are for $i = 0, 1, 2, 3$.

- (i) $e_0 = 0$ since a connected 0-manifold is a point;
- (ii) $e_1 = 0$ since a connected, closed 1-manifold is homeomorphic to S^1 ;
- (iii) $e_2 = 1$ since any orientable, connected, closed surface embeds into S^3 , and any non-orientable, connected, closed surface can be presented as a connected sum of an orientable, connected, closed surface and $\mathbb{RP}^2$ or $\mathbb{RP}^2 \# \mathbb{RP}^2$, hence each connected closed surface can be embedded into $S^3 \# \mathbb{RP}^3 \# \mathbb{RP}^3 \cong \mathbb{RP}^3 \# \mathbb{RP}^3$.

(iv) $e_3 = 2$ since each closed 3-manifold embeds into S^5 , due to Hirsch in the orientable case ([Hi], also [Ki]), and due to Rohlin and Wall independently in the non-orientable case ([Ro] [Wa], see [WZ] for a possibly more elementary proof, in the sense that only Dehn surgery is involved, of this Hirsch-Rohlin-Wall's Theorem); and Kawauchi showed that there does not exist a single oriented, connected, closed 4-manifold such that any connected, closed 3-manifold can be topologically flat embedded into it by constructing signature invariants from infinite cyclic coverings [Ka], and then Shiomi showed that Kawauchi's result is still true if the target 4-manifold is allowed to be non-orientable [Shi].

It is reasonable to guess that in general $e_n > 1$. The main result of the present paper claims that this is the case for “most” n , and indeed we prove a stronger version for those n 's.

Theorem 4.1. *If n is a composite number and is not a power of 2, then there does not exist a single connected, closed $(n + 1)$ -manifold W , such that any smooth, simply-connected, closed n -manifold M can be topologically flat embedded into W .*

We say that M can be *topologically flat* embedded into W if $M \times [0, 1]$ can be embedded into W .

Besides [Ka] and [Shi], recently there are several papers studying the embedding of a 3-manifold M into simply-connected 4-manifolds other than S^4 , say [EL], [Fa1], [Fa2], and indeed the targets in those papers are connected sums of $\mathbb{CP}^2$. This relates to finding uncountably many smooth structures on the 4-manifold $M \times \mathbb{R}$ (see [Di], [Fa1] and [Fa2]). Fang's observation (cf. [Fa2]) that there exists an obstruction to the embedding of 3-manifolds into a simply connected definite 4-manifold was the first inspiration of our proof for Theorem 4.1.

We expect that Theorem 4.1 still holds for all remaining positive integers $n \geq 5$. When $n = 3$, the only simply-connected, closed 3-manifold is S^3 if Poincaré Conjecture is valid. When $n = 4$, the situation is still not clear, and we have the following

Theorem 5.5. (a) *There exists an oriented, connected, closed 5-manifold W , such that for any simply-connected, closed 4-manifold M with indefinite intersection form, M can be topologically flat embedded into W .*

(b) *Any simply-connected, compact topological 4-manifold M with $\partial M \neq \emptyset$ admits a topologically flat embedding into $S^2 \tilde{\times} S^3$.*

Remark. There are indefinite 4-manifolds with arbitrarily large signatures. Theorem 5.5 (a) tells us that the signature of 4-manifolds cannot be an obstruction for our codimension 1 embeddings. Note that any oriented, connected, closed 4-manifold with non-zero signature does not bound a compact orientable

5-manifold. It follows that the first Betti number of W in (a) must be positive and that the condition $\partial M \neq \emptyset$ in (b) can not be removed.

Compared with Theorem 5.5 (b), it is worthy noting that there does not exist a single oriented, connected, closed 4-manifold such that any compact punctured 3-manifold can be topologically flat embedded into it (cf. [Ka]).

The organization of the paper and an outline of the proofs.

In Section 2, we first reduce the proof of Theorem 4.1 to the oriented case. Then for any fixed $(n+1)$ -manifold W , Proposition 2.1 claims that for each topologically flat codimension 1 embedding $M^n \rightarrow W$ and each factorization $n = pq$, there is a homological obstruction in terms of $H^{p+1}(W; \mathbb{R})$, $H^p(M; \mathbb{R})$, the q -multiple cup product on $H^p(M; \mathbb{R})$ and so on. Since n is not a prime number and is not a power of 2, we can assume that the factorization $n = pq$ has been chosen so that $p \geq 2$ and $q \geq 3$, and q is odd.

Let $(\wedge^q \mathbb{R}^m)^*$ (resp. $(\vee^q \mathbb{R}^m)^*$) be the vector space of all skew-symmetric (resp. symmetric) q -multilinear functions on the m -dimensional real vector space $\mathbb{R}^m$. Motivated by the homological obstruction in Proposition 2.1, in Section 3 we define when a function in $(\wedge^q \mathbb{R}^m)^*$ (resp. $(\vee^q \mathbb{R}^m)^*$) is special so that the q -multiple cup product on $H^p(M; \mathbb{R})$ in Proposition 2.1 is a special function. Then Proposition 3.1 claims that under certain circumstances, such special functions are contained in a proper closed subset of $(\wedge^q \mathbb{R}^m)^*$ (resp. $(\vee^q \mathbb{R}^m)^*$). The conditions that $q \geq 3$, and q is odd are used in proving Proposition 3.1.

In Section 4, we first choose a suitable rational non-special function provided by Proposition 3.1, and then Proposition 4.2 claims that there is a commutative graded algebra $A = \bigoplus_{i=0}^q A_{ip}$ over $\mathbb{Q}$ satisfying Poincaré duality so that the chosen non-special function is the q -multiple product on A_p . Next we verify that A can be realized as the rational cohomology ring of a simply-connected closed n -manifold M by invoking a celebrated result of Sullivan (the condition $p \geq 2$ is used here). Hence the q -multiple cup product on $H^p(M; \mathbb{R})$ is a non-special function, and therefore M cannot be topologically flat embedded into W by Proposition 2.1, which finishes the proof of Theorem 4.1.

Section 5 is devoted to the case $n = 4$, and we get Theorem 5.5 by explicit constructions and the classification theorem for simply-connected, closed 4-manifolds.

In the remaining of this section, We recall and fix needed notations and conventions in tensor algebra based on [Sh].

Prerequisites on tensor algebra. Let V be a vector space of finite dimension m over a field $\mathbb{F}$ of characteristic zero. The space $\otimes^p V$, consisting of contravariant tensors of order p , is defined for $p = 2, 3, \dots$ to be the tensor product $V \otimes V \otimes \dots \otimes V$ of p copies of V , and for $p = 0, 1$ to be $\mathbb{F}, V$. We

may identify $(\otimes^p V) \otimes (\otimes^q V)$ with $\otimes^{p+q} V$. Thus given contravariant tensors x, y of orders p, q we may form their product $x \otimes y$, a contravariant tensor of order $p + q$. Observe that the product of decomposable elements is given by $(x_1 \otimes \dots \otimes x_p) \otimes (y_1 \otimes \dots \otimes y_q) = x_1 \otimes \dots \otimes x_p \otimes y_1 \otimes \dots \otimes y_q$. Form the direct sum

$$\otimes V = \oplus_p (\otimes^p V) = \mathbb{F} \oplus V \oplus (\otimes^2 V) \oplus \dots$$

of all the spaces $\otimes^p V, p = 0, 1, 2, \dots$. If we extend the product operation $\otimes$ in the obvious bilinear way to general elements of the vector space $\otimes V$, we make $\otimes V$ into an associative algebra over $\mathbb{F}$, called the *tensor algebra* of the space V .

Let $N^p(V)$ denote the subspace of $\otimes^p V$ spanned by those tensors of the form $v_1 \otimes \dots \otimes v_p$, where $v_i = v_{i+1}$ for some $i = 1, 2, \dots, p-1$. For $p = 0$ or 1 , $N^p(V)$ is defined to be 0 . The p th *exterior power* of V is defined to be the quotient space $\wedge^p V = \otimes^p V / N^p(V)$. Observe that $\wedge^0 V = \mathbb{F}$, $\wedge^1 V = V$. The canonical projection $\pi_p : \otimes^p V \rightarrow \wedge^p V$ is defined by $\pi_p(x) = x + N^p(V)$ for $x \in \otimes^p V$. For $v_1, \dots, v_p \in V$, we will write $\pi_p(v_1 \otimes \dots \otimes v_p)$ as $v_1 \wedge \dots \wedge v_p$. There is a product operation $\wedge : (\wedge^p V) \times (\wedge^q V) \rightarrow \wedge^{p+q} V$ defined as follows. If $x \in \otimes^p V, y \in \otimes^q V$, then the *exterior product* of $\pi_p(x)$ and $\pi_q(y)$ is defined by $\pi_p(x) \wedge \pi_q(y) = \pi_{p+q}(x \otimes y)$. Observe that the exterior product of decomposable elements is given by $(x_1 \wedge \dots \wedge x_p) \wedge (y_1 \wedge \dots \wedge y_q) = x_1 \wedge \dots \wedge x_p \wedge y_1 \wedge \dots \wedge y_q$. If we extend $\wedge$ in the obvious bilinear way to general elements of the vector space

$$\wedge V = \oplus_p (\wedge^p V) = \mathbb{F} \oplus V \oplus \wedge^2 V \oplus \dots \oplus \wedge^m V,$$

we make $\wedge V$ into an algebra over $\mathbb{F}$, called the *exterior algebra* on V .

Let S_p denote the symmetric group on the p symbols $\{1, 2, \dots, p\}$. Each permutation $\sigma \in S_p$ gives rise to a linear operator, still denoted by σ , on $\otimes^p V$ defined by its effect

$$\sigma(v_1 \otimes \dots \otimes v_p) = v_{\sigma^{-1}(1)} \otimes \dots \otimes v_{\sigma^{-1}(p)}$$

upon the decomposable elements of $\otimes^p V$. Let $K^p(V)$ denote the subspace of $\otimes^p V$ spanned by those tensors of the form $u - \sigma u$ for some $u \in \otimes^p V$ and some $\sigma \in S_p$. If $p = 0$ or 1 we define $K^p(V)$ to be 0 . The p th *symmetric power* of V is defined to be the quotient space $\vee^p V = \otimes^p V / K^p(V)$. Observe that $\vee^0 V = \mathbb{F}$, $\vee^1 V = V$. The canonical projection $\psi_p : \otimes^p V \rightarrow \vee^p V$ is defined by $\psi_p(x) = x + K^p(V)$ for $x \in \otimes^p V$. Then exactly parallel to the description in the preceding paragraph (simply changing π_p to ψ_p and $\wedge$ to $\vee$), we see that there is a product operation $\vee : (\vee^p V) \times (\vee^q V) \rightarrow \vee^{p+q} V$ which can be extended in the obvious bilinear way to general elements of the vector space

$$\vee V = \oplus_p (\vee^p V) = \mathbb{F} \oplus V \oplus \vee^2 V \oplus \dots$$

to make $\vee V$ into an infinite-dimensional algebra over $\mathbb{F}$, called the *symmetric algebra* on V .

Let $\times^p V$ denote the Cartesian product $V \times V \times \dots \times V$ of p copies of V . Each multilinear function $F : \times^p V \rightarrow \mathbb{F}$ determines a unique linear function $L : \otimes^p V \rightarrow \mathbb{F}$ such that for all $v_1, \dots, v_p \in V$, $L(v_1 \otimes \dots \otimes v_p) = F(v_1, \dots, v_p)$. The correspondence $F \leftrightarrow L$ establishes a natural isomorphism between the vector space $L(\times^p V; \mathbb{F})$ of all multilinear functions and the dual space $(\otimes^p V)^*$ of $\otimes^p V$. For any skew-symmetric (resp. symmetric) multilinear function $F : \times^p V \rightarrow \mathbb{F}$, there exists a unique linear function $L_s : \wedge^p V$ (resp. $\vee^p V$) $\rightarrow \mathbb{F}$ such that for all $v_1, \dots, v_p \in V$, $L_s(v_1 \wedge \dots \wedge v_p) = F(v_1, \dots, v_p)$ (resp. $L_s(v_1 \vee \dots \vee v_p) = F(v_1, \dots, v_p)$). The correspondence $F \leftrightarrow L_s$ establishes a natural isomorphism between the vector space $\text{Sk}(\times^p V; \mathbb{F})$ (resp. $\text{Sym}(\times^p V; \mathbb{F})$) of all skew-symmetric (resp. symmetric) multilinear functions and the dual space $(\wedge^p V)^*$ (resp. $(\vee^p V)^*$) of $\wedge^p V$ (resp. $\vee^p V$). Thus we may consider a skew-symmetric (resp. symmetric) multilinear function $F : \times^p V \rightarrow \mathbb{F}$ as an element of $(\wedge^p V)^*$ (resp. $(\vee^p V)^*$), and vice versa.

Let $\{e_1, \dots, e_m\}$ be the standard basis for the real vector space $\mathbb{R}^m$, i.e. $e_i = (0, \dots, 0, 1, 0, \dots, 0)$ has components 0 except for its i -th component, which is equal to 1. Note that $(\wedge^q \mathbb{R}^m)^*$ (resp. $(\vee^q \mathbb{R}^m)^*$) has dimension $\binom{m}{q}$ (resp. $\binom{m+q-1}{q}$). Each element $F \in (\wedge^q \mathbb{R}^m)^*$ (resp. $(\vee^q \mathbb{R}^m)^*$) is determined by $\binom{m}{q}$ (resp. $\binom{m+q-1}{q}$) real numbers $(F_{i_1 \dots i_q})$, where $F_{i_1 \dots i_q} = F(e_{i_1}, \dots, e_{i_q})$ for $1 \leq i_1 < \dots < i_q \leq m$ (resp. $1 \leq i_1 \leq \dots \leq i_q \leq m$). We call $F_{i_1 \dots i_q}$ the $(i_1, \dots, i_q)$ component of F .

2 Reduction to the oriented case and homological obstructions for the embedding

If any smooth, simply-connected, closed n -manifold M can be topologically flat embedded into a non-orientable $(n+1)$ -manifold W , then M can also be topologically flat embedded into the orientable double cover of W because M is simply-connected. It follows that Theorem 4.1 is equivalent to the following

Theorem 4.1*. *If n is a composite number and is not a power of 2, then there does not exist a single oriented, connected, closed $(n+1)$ -manifold W , such that any smooth, simply-connected, closed n -manifold M can be topologically flat embedded into W .*

From now on all manifolds appearing in this paper are considered to be

oriented and connected. $\beta_i(X)$ denotes the i -th Betti number of a compact manifold X .

The following proposition sets up a property on the cohomology ring of n -manifold M if it can be embedded into the $(n+1)$ -manifold W .

Proposition 2.1. *Let M and W be closed, connected, oriented n - and $(n+1)$ -dimensional manifolds respectively.*

If M topologically flat embeds in W , then for any integer factorization $n = pq$, where $p, q > 0$, there exists a subspace V of $H^p(M; \mathbb{R})$ and a linear transformation $\varphi : V \rightarrow H^p(M; \mathbb{R})$ such that

- (i) φ has no fixed non-zero vectors, i.e. for any non-zero $x \in V$, $\varphi(x) \neq x$,
- (ii) $\dim V \geq \frac{1}{2}(\beta_p(M) - \beta_{p+1}(W))$,
- (iii) for any $x_1, \dots, x_q \in V$, $x_1 \cup \dots \cup x_q = \varphi(x_1) \cup \dots \cup \varphi(x_q)$.

Proof. All homology and cohomology groups in this proof are with coefficients in $\mathbb{R}$. We divide the proof into two cases.

Case 1. Suppose $W \setminus M$ is not connected. Denote the closures of the two components of $W \setminus M$ by W_1 and W_2 . By Mayer-Vietoris sequence, we know that

$$H^p(W_1) \oplus H^p(W_2) \xrightarrow{f} H^p(M) \xrightarrow{\delta^*} H^{p+1}(W)$$

is exact. Here the first map is given by $f(w_1, w_2) = i_1^*(w_1) - i_2^*(w_2)$ for $w_1 \in H^p(W_1), w_2 \in H^p(W_2)$, where i_1 and i_2 are inclusions $M \hookrightarrow W_1$ and $M \hookrightarrow W_2$ respectively. Clearly

$$\dim \operatorname{im}(f) = \dim \ker(\delta^*) \geq \dim H^p(M) - \dim H^{p+1}(W) = \beta_p(M) - \beta_{p+1}(W).$$

Since $\dim \operatorname{im}(i_1^*) + \dim \operatorname{im}(i_2^*) \geq \dim \operatorname{im}(f)$, we may set $V = \operatorname{im}(i_1^*)$ and assume that $\dim V \geq \frac{1}{2}(\beta_p(M) - \beta_{p+1}(W))$.

For any $x_1, \dots, x_q \in V$, there exist $y_1, \dots, y_q \in W_1$ such that $x_i = i_1^*(y_i)$, $1 \leq i \leq q$. Since $M = \partial W_1$, we have $i_{1*}[M] = 0 \in H_n(W_1)$, where $[M]$ denotes the fundamental class of M . So

$$\begin{aligned} \langle x_1 \cup \dots \cup x_q, [M] \rangle &= \langle i_1^*(y_1 \cup \dots \cup y_q), [M] \rangle \\ &= \langle y_1 \cup \dots \cup y_q, i_{1*}[M] \rangle = 0. \end{aligned}$$

Therefore for any $x_1, \dots, x_q \in V$, $x_1 \cup \dots \cup x_q = 0$.

The result is thus proved by letting φ to be the zero map.

Case 2. Suppose $W \setminus M$ is connected. We identify a neighborhood W_1 of M in W with $M \times I$ and let $W_2 = W \setminus \operatorname{int}(W_1)$, where $\operatorname{int}(W_1)$ denotes the interior of W_1 . Clearly, $W_1 \cap W_2 = (M \times \{0\}) \cup (M \times \{1\})$.

Still by Mayer-Vietoris sequence, we get the exact sequence

$$H^p(W_1) \oplus H^p(W_2) \rightarrow H^p(W_1 \cap W_2) \rightarrow H^{p+1}(W)$$

and it is reduced to

$$H^p(M) \oplus H^p(W_2) \xrightarrow{f} H^p(M) \oplus H^p(M) \xrightarrow{\delta^*} H^{p+1}(W).$$

Now f can be written as

$$f(x, w) = (x - i_0^*(w), x - i_1^*(w)),$$

where i_0 and i_1 are inclusions $M \times \{0\} \hookrightarrow W_2$ and $M \times \{1\} \hookrightarrow W_2$ respectively.

By the exactness,

$$\dim \operatorname{im}(f) = \dim \ker(\delta^*) \geq \dim(H^p(M) \oplus H^p(M)) - \dim H^{p+1}(W) = 2\beta_p(M) - \beta_{p+1}(W).$$

Let $\Delta = \{(x, x) | x \in H^p(M)\} \subset H^p(M) \oplus H^p(M)$ and $U = \{(i_0^*(w), i_1^*(w)) | w \in H^p(W_2)\}$, then $\Delta + U = \operatorname{im}(f)$ and therefore

$$\dim(\Delta + U) = \dim \operatorname{im}(f) \geq 2\beta_p(M) - \beta_{p+1}(W).$$

Since $\dim \Delta = \beta_p(M)$, there is a subspace S of U with $\dim S \geq \beta_p(M) - \beta_{p+1}(W)$ such that $S \cap \Delta = \{(0, 0)\}$. Let $\pi_1, \pi_2 : H^p(M) \oplus H^p(M) \rightarrow H^p(M)$ be the projections of $H^p(M) \oplus H^p(M)$ onto its first and second factors, respectively. Since $\dim \pi_1(S) + \dim \pi_2(S) \geq \dim S$, without loss of generality, we may assume that

$$\dim \pi_1(S) \geq \frac{1}{2} \dim S \geq \frac{1}{2}(\beta_p(M) - \beta_{p+1}(W)).$$

It follows from $\partial W_2 = -(M \times \{0\}) \cup (M \times \{1\})$ that $i_{0*}[M] = i_{1*}[M]$. Thus for any $w_1, \dots, w_q \in H^p(W_2)$, $\langle w_1 \cup \dots \cup w_q, i_{0*}[M] - i_{1*}[M] \rangle = 0$, and this implies

$$\langle i_0^*(w_1) \cup \dots \cup i_0^*(w_q), [M] \rangle = \langle i_1^*(w_1) \cup \dots \cup i_1^*(w_q), [M] \rangle.$$

In particular, if $(u_1, v_1), \dots, (u_q, v_q) \in S$, then

$$u_1 \cup \dots \cup u_q = v_1 \cup \dots \cup v_q. \quad (1)$$

Let $\{y_1, \dots, y_r\}$ be a basis for $\pi_1(S)$. We can find $z_1, \dots, z_r \in H^p(M)$ such that $(y_j, z_j) \in S$, $1 \leq j \leq r$. Then there is a unique linear transformation

$$\varphi : \pi_1(S) \rightarrow H^p(M)$$

such that $\varphi(y_j) = z_j$, $1 \leq j \leq r$.

One can write any $y \in \pi_1(S)$ as $y = \sum_j c_j y_j$, then $\varphi(y) = \sum_j c_j z_j$. Thus $(y, \varphi(y)) \in S$. Since $S \cap \Delta = \{(0, 0)\}$, if $y \neq 0$, then $\varphi(y) \neq y$. Now set $V = \pi_1(S)$ and one gets the desired result from equation (1). $\square$

3 Special multilinear functions

Proposition 2.1 motivates the following definition.

Definition. Let F be an element of $(\wedge^q \mathbb{R}^m)^*$ (resp. $(\vee^q \mathbb{R}^m)^*$). We say that F is *special* if there exist a subspace U of $\mathbb{R}^m$ with $\dim U \geq \frac{m}{3}$ and a linear map $\varphi : U \rightarrow \mathbb{R}^m$ with no fixed non-zero vectors (i.e. for any non-zero $x \in U$, $\varphi(x) \neq x$) such that for all $x_1, \dots, x_q \in U$,

$$F(x_1, \dots, x_q) = F(\varphi(x_1), \dots, \varphi(x_q)).$$

In the above equation, we consider F as a skew-symmetric (resp. symmetric) multilinear function (cf. Section 1).

In this section, we want to show

Proposition 3.1. *Suppose $q \geq 3$ is an odd integer.*

(a) *If m is sufficiently large, then there exists a proper closed subset X_m of $(\wedge^q \mathbb{R}^m)^*$ such that if $F \in (\wedge^q \mathbb{R}^m)^*$ is special, then $F \in X_m$.*

(b) *The result in (a) is still true if we substitute $(\wedge^q \mathbb{R}^m)^*$ by $(\vee^q \mathbb{R}^m)^*$.*

Remark. If $q \geq 2$ is even, then each $F \in (\wedge^q \mathbb{R}^m)^*$ (resp. $F \in (\vee^q \mathbb{R}^m)^*$) is special. We may simply take $U = \mathbb{R}^m$ and $\varphi : \mathbb{R}^m \rightarrow \mathbb{R}^m$ to be the map sending x to $-x$ for all $x \in \mathbb{R}^m$.

Before proving Proposition 3.1, we shall discuss some useful lemmas. The first lemma describes the real Jordan canonical form.

Lemma 3.2. *Any real square matrix is similar to a block diagonal matrix, where each block takes one of the following two forms*

$$J_k(a) = \begin{bmatrix} a & 1 & & \\ & a & 1 & \\ & & \ddots & \ddots \\ & & & \ddots & 1 \\ & & & & a \end{bmatrix}, C_l(a, b) = \begin{bmatrix} a & b & 1 & 0 & & \\ -b & a & 0 & 1 & & \\ & a & b & \ddots & & \\ -b & a & & \ddots & & \\ & & \ddots & & 1 & 0 \\ & & & \ddots & 0 & 1 \\ & & & & a & b \\ & & & & -b & a \end{bmatrix}.$$

The sizes of these two square matrices are k and $2l$ respectively, where k and l are positive integers, a and $b \neq 0$ are real numbers.

Proof. This is Theorem 3.4.5 in [HJ]. □

Now we sort the blocks described in this lemma into 6 types for later use.

Type 1: $J_1(1)$, **Type 2:** $J_1(-1)$, **Type 3:** $J_1(a), |a| > 1$,
Type 4: $J_1(a), |a| < 1$, **Type 5:** $J_k(a), k > 1$, **Type 6:** $C_l(a, b)$.

Lemma 3.3. *Let V be a subspace of $\mathbb{R}^m$ with $\dim V = l$. Suppose that $\{v_1, \dots, v_l\}$ is a basis for V , and that $\{v_1, \dots, v_l, \dots, v_m\}$ is a basis for $\mathbb{R}^m$.*

*Let $\varphi : V \rightarrow \mathbb{R}^m$ be a linear map with no fixed non-zero vectors such that the matrix of φ with respect to the above bases is $\begin{bmatrix} J \\ * \end{bmatrix}$, where J is an $l \times l$ block diagonal matrix and all the blocks are of Type i_0 , $i_0 = 1, 5$ or 6 . Then there exists a subspace V' of V such that $\dim V' \geq \frac{l}{3}$, $\varphi(V') \cap V' = \{0\}$ and φ is injective on V' .*

Proof. First suppose that J only contains Type 1 blocks. Let $W = \text{span}\{v_{l+1}, \dots, v_m\}$, then $\mathbb{R}^m = V \oplus W$. For any $v \in V$, since $J = I$, we have decomposition

$$\varphi(v) = v + w, v \in V, w \in W.$$

Thus φ is injective. Suppose that $x \in \varphi(V) \cap V$, then $x \in V$ and there exists $v \in V$ such that $\varphi(v) = x$. Now the decomposition of $\varphi(v)$ under $V \oplus W$ equals both $x + 0$ and $v + w$, so $x = v$, $\varphi(v) = v$. Since φ only fixes the zero vector, we have $v = 0$. Therefore $\varphi(V) \cap V = \{0\}$, and we may simply choose $V' = V$.

Suppose now that J only contains Type 5 or Type 6 blocks. For a block in J , let $\{w_1, \dots, w_k\}$ be the subset of the basis $\{v_1, \dots, v_l\}$ that corresponds to that block. Note that $k \geq 2$. Set $W = \text{span}\{w_1, \dots, w_k\}$, $W_0 = \text{span}\{w_j \mid j \text{ even}\}$ and $W_1 = \text{span}\{w_j \mid j \text{ odd}\}$. Clearly, $W = W_0 \oplus W_1$ and $\dim W_0 \geq \frac{k}{3}$. Denote the projection of $\mathbb{R}^m$ onto W_1 by p_1 . (This projection is well-defined as there is a natural decomposition $\mathbb{R}^m = W \oplus W'$, where $W' = \text{span}(\{v_1, \dots, v_m\} \setminus \{w_1, \dots, w_k\})$.) If the block is $C_{\frac{k}{2}}(a, b)$, then

$$p_1 \varphi(w_{2i}) = bw_{2i-1}, \quad 1 \leq i \leq \frac{k}{2}, \quad (2)$$

and if the block is $J_k(a)$, then

$$p_1 \varphi(w_{2i}) = w_{2i-1}, \quad 1 \leq i \leq \left\lfloor \frac{k}{2} \right\rfloor, \quad (3)$$

where $\left\lfloor \frac{k}{2} \right\rfloor$ is the greatest integer $\leq \frac{k}{2}$. Note that $b \neq 0$. In any case, $p_1 \varphi(x) = 0$ holds for $x \in W_0$ only if $x = 0$. Thus φ is injective on W_0 . By equations (2) and (3), $\varphi(W_0) \cap W_0 = \{0\}$. Now we may choose V' to be the direct sum of these W_0 's of all blocks in J . $\square$

The next lemma needs to use the transcendence degree of a field extension. A good reference is [La].

Lemma 3.4. *Let $g_1(x_1, \dots, x_r), \dots, g_s(x_1, \dots, x_r)$ be rational functions in r variables $x_1, \dots, x_r$ with coefficients in $\mathbb{R}$. If $s > r$, then there exists a non-zero polynomial P in s variables such that*

$$P(g_1(x_1, \dots, x_r), \dots, g_s(x_1, \dots, x_r)) = 0.$$

Proof. Let K be the field of fractions of the polynomial ring $\mathbb{R}[x_1, \dots, x_r]$. Clearly, $\{x_1, \dots, x_r\}$ is a transcendence base of K over $\mathbb{R}$. Since $s > r$, $\{g_1, \dots, g_s\}$ is not algebraically independent over $\mathbb{R}$ (cf. [La], Chapter VIII, Theorem 1.1). Hence there exists a non-zero polynomial P in s variables with coefficients in $\mathbb{R}$ such that $P(g_1, \dots, g_s) = 0$. $\square$

Proof of Proposition 3.1.

(a) Suppose that $F_0 \in (\wedge^q \mathbb{R}^m)^*$ is special. Thus there exist a subspace U of $\mathbb{R}^m$ with $\dim U = k \geq \frac{m}{3}$ and a linear map $\varphi : U \rightarrow \mathbb{R}^m$ with no fixed non-zero vectors such that for all $x_1, \dots, x_q \in U$,

$$F_0(x_1, \dots, x_q) = F_0(\varphi(x_1), \dots, \varphi(x_q)).$$

We will use the above equation to get expressions for coordinates of F_0 under standard basis for $(\wedge^q \mathbb{R}^m)^*$. The number of variables used will be less than $\dim(\wedge^q \mathbb{R}^m)^*$ when m is large enough. Hence all special F_0 lie in the zero set of a single non-zero polynomial, which is a proper closed subset of $(\wedge^q \mathbb{R}^m)^*$.

Choose vectors $u_{k+1}, \dots, u_m$ such that

$$U \oplus \text{span}\{u_{k+1}, \dots, u_m\} = \mathbb{R}^m.$$

Let $\{u_1, \dots, u_k\}$ be a basis for U , then $\{u_1, \dots, u_k, u_{k+1}, \dots, u_m\}$ is a basis for $\mathbb{R}^m$. Denote the matrix of φ with respect to these two bases by $\begin{bmatrix} J \\ * \end{bmatrix}$, where J is a $k \times k$ square matrix. A suitable choice of $\{u_1, \dots, u_k\}$ can make J in real Jordan canonical form as in Lemma 3.2. The Jordan blocks have 6 types. For some i_0 , the total size of blocks of Type i_0 in J should be at least $\frac{k}{6} \geq \frac{m}{18}$. Denote this total size by l . We change the order of $u_1, \dots, u_k$ so that in the block expression for J , Type i_0 blocks appear first. From now on, vectors $u_1, \dots, u_k, \dots, u_m$ are fixed.

Case 1. Suppose that $i_0 = 1, 5$ or 6 . Then by Lemma 3.3, we may find a subspace U' of $\text{span}\{u_1, \dots, u_l\}$ with $k' = \dim U' \geq \frac{l}{3} \geq \frac{m}{54}$, $\varphi(U') \cap U' = \{0\}$ and φ is injective on U' . Let m_1 denote $[\frac{m}{54}]$, the greatest integer $\leq \frac{m}{54}$. Let $\{v_1, \dots, v_{k'}\}$ be a basis of U' . Then $\{v_1, \dots, v_{k'}, \varphi(v_1), \dots, \varphi(v_{k'})\}$ is linearly independent and there exists an isomorphism $\tau_1 : \mathbb{R}^m \rightarrow \mathbb{R}^m$ sending v_i to e_i , $\varphi(v_i)$ to e_{m-m_1+i} for $1 \leq i \leq m_1$. (cf. section 1 for definition of e_i .)

Let $L(\mathbb{R}^m, \mathbb{R}^m)$ be the vector space consisting of all linear maps from $\mathbb{R}^m$ to $\mathbb{R}^m$. $L(\mathbb{R}^m, \mathbb{R}^m)$ has dimension m^2 . Each element $\tau \in L(\mathbb{R}^m, \mathbb{R}^m)$ is determined

by m^2 numbers (τ_{ji}) , where $\tau(e_i) = \sum_{j=1}^m \tau_{ji} e_j$ for $1 \leq i \leq m$. For each $\tau \in L(\mathbb{R}^m, \mathbb{R}^m)$, τ induces a linear map $\tau^* : (\wedge^q \mathbb{R}^m)^* \rightarrow (\wedge^q \mathbb{R}^m)^*$ defined by

$$(\tau^* F)(x_1, \dots, x_q) = F(\tau(x_1), \dots, \tau(x_q))$$

for $F \in (\wedge^q \mathbb{R}^m)^*$ and $x_1, \dots, x_q \in \mathbb{R}^m$.

Let $R = \{(i_1, \dots, i_q) | 1 \leq i_1 < \dots < i_q \leq m\}$, $R_1 = \{(i_1, \dots, i_q) | 1 \leq i_1 < \dots < i_q \leq m_1\}$ and $R'_1 = R \setminus R_1$.

Now consider $(\tau_1^{-1})^* F_0$. Note that for $(i_1, \dots, i_q) \in R_1$, we have

$$\begin{aligned} ((\tau_1^{-1})^* F_0)_{i_1 \dots i_q} &= ((\tau_1^{-1})^* F_0)(e_{i_1}, \dots, e_{i_q}) \\ &= F_0(v_{i_1}, \dots, v_{i_q}) = F_0(\varphi(v_{i_1}), \dots, \varphi(v_{i_q})) \\ &= ((\tau_1^{-1})^* F_0)(e_{m-m_1+i_1}, \dots, e_{m-m_1+i_q}) \\ &= ((\tau_1^{-1})^* F_0)_{(m-m_1+i_1) \dots (m-m_1+i_q)}. \end{aligned}$$

Let T denote the subspace of $(\wedge^q \mathbb{R}^m)^*$ consisting of all $F \in (\wedge^q \mathbb{R}^m)^*$ such that $F_{i_1 \dots i_q} = F_{(m-m_1+i_1) \dots (m-m_1+i_q)}$ for all $(i_1, \dots, i_q) \in R_1$. Note that $(\tau_1^{-1})^* F_0 \in T$. As each element $F \in T$ is determined by $\binom{m}{q} - \binom{m_1}{q}$ numbers $(F_{i_1 \dots i_q})_{(i_1, \dots, i_q) \in R'_1}$. Thus, T is a vector space of dimension $\binom{m}{q} - \binom{m_1}{q}$.

Define $\Psi_1 : L(\mathbb{R}^m, \mathbb{R}^m) \times T \rightarrow (\wedge^q \mathbb{R}^m)^*$ by $\Psi_1(\tau, F) = \tau^* F$ for $\tau \in L(\mathbb{R}^m, \mathbb{R}^m)$ and $F \in T$. Since $(\tau_1^{-1})^* F_0 \in T$, we have $F_0 \in \text{im}(\Psi_1)$. Observe that for each $(i_1, \dots, i_q) \in R$, the $(i_1, \dots, i_q)$ component of $\Psi_1(\tau, F)$, $(\tau^* F)_{i_1 \dots i_q}$, is a polynomial of τ_{ji} 's and $F_{j_1 \dots j_q}$'s, where $1 \leq i, j \leq m$ and $(j_1, \dots, j_q) \in R'_1$. Identify $(\wedge^q \mathbb{R}^m)^*$ with $\mathbb{R}^{a_0}$ through standard basis $\{e_1, \dots, e_m\}$, where $a_0 = \binom{m}{q}$. Note that since $q \geq 3$, if m is sufficiently large, then $m^2 + \binom{m}{q} - \binom{m_1}{q} < \binom{m}{q}$. Thus by Lemma 3.4, we can find a non-zero polynomial P_1 such that

$$\text{im}(\Psi_1) \subset \{x \in \mathbb{R}^{a_0} | P_1(x) = 0\}.$$

Case 2. Suppose that $i_0 = 2, 3$ or 4 . Let $\tau_2 : \mathbb{R}^m \rightarrow \mathbb{R}^m$ be the isomorphism sending u_i to e_i for $1 \leq i \leq m$. Let m_2 denote $\lceil \frac{m}{18} \rceil$, then $m_2 \leq l$ and for $1 \leq i \leq m_2$,

$$(\tau_2 \varphi \tau_2^{-1})(e_i) = \lambda_i e_i + \sum_{j=m_2+1}^m d_{ji} e_j,$$

for some λ_i and d_{ji} 's. Let $R_2 = \{(i_1, \dots, i_q) | 1 \leq i_1 < \dots < i_q \leq m_2\}$ and $R'_2 = R \setminus R_2$. Note that for $(i_1, \dots, i_q) \in R_2$, we have $\lambda_{i_1} \dots \lambda_{i_q} \neq 1$. (If $i_0 = 2$,

then $\lambda_{i_1} \dots \lambda_{i_q} = -1$ since q is odd; if $i_0 = 3$, then $|\lambda_{i_1} \dots \lambda_{i_q}| > 1$; if $i_0 = 4$, then $|\lambda_{i_1} \dots \lambda_{i_q}| < 1$.) Denote $(\tau_2^{-1})^* F_0$ by G . Then for $(i_1, \dots, i_q) \in R_2$,

$$F_0(u_{i_1}, \dots, u_{i_q}) = F_0(\varphi(u_{i_1}), \dots, \varphi(u_{i_q})),$$

implies

$$\begin{aligned} G_{i_1 \dots i_q} &= G(e_{i_1}, \dots, e_{i_q}) = G((\tau_2 \varphi \tau_2^{-1})(e_{i_1}), \dots, (\tau_2 \varphi \tau_2^{-1})(e_{i_q})) \\ &= \lambda_{i_1} \dots \lambda_{i_q} G_{i_1 \dots i_q} + \sum_{(j_1, \dots, j_q) \in R'_2} c_{i_1 \dots i_q, j_1 \dots j_q} G_{j_1 \dots j_q}, \end{aligned}$$

or equivalently,

$$G_{i_1 \dots i_q} = \frac{\sum_{(j_1, \dots, j_q) \in R'_2} c_{i_1 \dots i_q, j_1 \dots j_q} G_{j_1 \dots j_q}}{1 - \lambda_{i_1} \dots \lambda_{i_q}}, \quad (4)$$

where each coefficient $c_{i_1 \dots i_q, j_1 \dots j_q}$ is a polynomial of λ_i 's and d_{ji} 's. Hence for any $(i_1, \dots, i_q) \in R_2$, $G_{i_1 \dots i_q}$ can be expressed as a rational function of λ_i 's, d_{ji} 's and $G_{j_1 \dots j_q}$'s, where $1 \leq i \leq m_2, m_2 + 1 \leq j \leq m$ and $(j_1, \dots, j_q) \in R'_2$.

Let Δ denote the open subset of $\mathbb{R}^{m_2}$ consisting of all m_2 -tuples $(\lambda_1, \dots, \lambda_{m_2}) \in \mathbb{R}^{m_2}$ such that for each $(i_1, \dots, i_q) \in R_2$, $\lambda_{i_1} \dots \lambda_{i_q} \neq 1$. Let $M_{(m-m_2) \times m_2}(\mathbb{R})$ denote the vector space consisting of all $(m - m_2) \times m_2$ matrices with entries in $\mathbb{R}$. Denote $\binom{m}{q} - \binom{m_2}{q}$ by b_0 .

Define $\Psi_2 : L(\mathbb{R}^m, \mathbb{R}^m) \times \Delta \times M_{(m-m_2) \times m_2}(\mathbb{R}) \times \mathbb{R}^{b_0} \rightarrow (\wedge^q \mathbb{R}^m)^*$ as follows: Let $\tau \in L(\mathbb{R}^m, \mathbb{R}^m)$, $\lambda = (\lambda_1, \dots, \lambda_{m_2}) \in \Delta$, $d = (d_{ji})_{m_2+1 \leq j \leq m, 1 \leq i \leq m_2} \in M_{(m-m_2) \times m_2}(\mathbb{R})$, $f = (f_{j_1 \dots j_q})_{(j_1, \dots, j_q) \in R'_2} \in \mathbb{R}^{b_0}$. Then there is a unique $F \in (\wedge^q \mathbb{R}^m)^*$ such that for each $(j_1, \dots, j_q) \in R'_2$, $F_{j_1 \dots j_q} = f_{j_1 \dots j_q}$ and for each $(i_1, \dots, i_q) \in R_2$, F satisfies equation (4) (changing G to F in that equation). Define $\Psi_2(\tau, \lambda, d, f)$ to be $\tau^* F$. Note that $F_0 \in \text{im}(\Psi_2)$. Observe that for each $(i_1, \dots, i_q) \in R$, the $(i_1, \dots, i_q)$ component of $\Psi_2(\tau, \lambda, d, f)$ is a rational function of τ_{ji} 's, λ_i 's, d_{ji} 's and $f_{j_1 \dots j_q}$'s. Identify $(\wedge^q \mathbb{R}^m)^*$ with $\mathbb{R}^{a_0}$, where $a_0 = \binom{m}{q}$. Note that since $q \geq 3$, if m is sufficiently large, then

$m^2 + m_2 + (m - m_2)m_2 + \binom{m}{q} - \binom{m_2}{q} < \binom{m}{q}$. Thus by Lemma 3.4, we can find a non-zero polynomial P_2 such that

$$\text{im}(\Psi_2) \subset \{x \in \mathbb{R}^{a_0} | P_2(x) = 0\}.$$

For sufficiently large m , let $P = P_1 P_2$ and $X_m = \{x \in \mathbb{R}^{a_0} | P(x) = 0\}$. X_m is obviously closed. It cannot be the whole $\mathbb{R}^{a_0}$ since P is a non-zero polynomial. Thus X_m is a proper closed subset of $(\wedge^q \mathbb{R}^m)^*$. If $F \in (\wedge^q \mathbb{R}^m)^*$ is special, then $F \in \text{im}(\Psi_1) \cup \text{im}(\Psi_2) \subset X_m$, as desired.

(b) We only need to substitute $(\wedge^q \mathbb{R}^m)^*$ by $(\vee^q \mathbb{R}^m)^*$, substitute $\binom{m_i}{q}$ by $\binom{m_i + q - 1}{q}$ where $i = \emptyset, 1$ or 2 and change the restriction $1 \leq i_1 < \dots < i_q \leq m$ to $1 \leq i_1 \leq \dots \leq i_q \leq m$ in the definition of R , R_1 and R_2 . $\square$

4 Main Theorem

In this section we prove the main theorem.

Theorem 4.1. *If n is a composite number and is not a power of 2, then there does not exist a single connected, closed $(n+1)$ -manifold W , such that any smooth, simply-connected, closed n -manifold M can be topologically flat embedded into W .*

We only need to prove

Theorem 4.1*. *If n is a composite number and is not a power of 2, then there does not exist a single oriented, connected, closed $(n+1)$ -manifold W , such that any smooth, simply-connected, closed n -manifold M can be topologically flat embedded into W .*

Definition. A *graded algebra* over $\mathbb{Q}$ is an algebra A over $\mathbb{Q}$ which as a vector space can be expressed as a direct sum

$$A = \bigoplus_{i=0}^{\infty} A_i,$$

and such that the bilinear multiplication $\cup : A \times A \rightarrow A$ maps $A_r \times A_s$ into A_{r+s} for all nonnegative integers r, s . A is *associative* if $(a \cup b) \cup c = a \cup (b \cup c)$ for all $a, b, c \in A$, and is *commutative* if for all $a \in A_r, b \in A_s (r, s \geq 0)$, $a \cup b = (-1)^{rs}(b \cup a)$. We always assume that A is associative, $A_0 = \mathbb{Q}$ and $1 \in \mathbb{Q} = A_0$ is a unit element of the algebra A .

Note that the rational cohomology ring $H^*(X; \mathbb{Q})$ of a path-connected topological space X is a commutative graded algebra over $\mathbb{Q}$ (cf. [Sp], pp. 263-264).

Definition. Let

$$A = \bigoplus_{i=0}^{\infty} A_i$$

be a commutative graded algebra over $\mathbb{Q}$. Assume that there exists a positive integer n such that $A_n \cong \mathbb{Q}$, $A_i = 0$ for $i > n$ and A_i is finite dimensional for $i < n$. Let φ_i be the map from A_i to $\text{Hom}(A_{n-i}, A_n)$ defined by

$$\varphi_i(u) : v \mapsto u \cup v, \quad \forall u \in A_i, v \in A_{n-i}.$$

We say that A satisfies *Poincaré duality* if φ_i is an isomorphism from A_i to $\text{Hom}(A_{n-i}, A_n)$ for any $0 \leq i \leq n$.

Proposition 4.2. *Suppose that p is a positive integer, $q \geq 3$ is an odd integer. F is in $(\wedge^q V)^*$ or $(\vee^q V)^*$ according to p is odd or even, where V is a finite dimensional vector space over $\mathbb{Q}$.*

There exists a commutative graded algebra

$$A = \bigoplus_{i=0}^{pq} A_i$$

satisfying Poincaré duality such that

- (i) $A_p = V$, $A_{pq} = \mathbb{Q}$, $A_i = 0$ if $i > pq$ and $A_i \neq 0$ only if i is a multiple of p ,
- (ii) for all $v_1, \dots, v_q \in A_p = V$, $F(v_1, \dots, v_q) = v_1 \cup \dots \cup v_q$.

Proof. First assume that p is odd. Let $q = 2r + 1$. We construct A by setting

$$A_i = \begin{cases} \wedge^s V & \text{if } i = sp \text{ and } 0 \leq s \leq r, \\ (\wedge^{2r+1-s} V)^* & \text{if } i = sp \text{ and } r+1 \leq s \leq 2r+1, \\ 0 & \text{otherwise} \end{cases}$$

Now suppose $u \in A_{sp}$, $v \in A_{tp}$ ($0 \leq s, t \leq 2r+1$). We define $u \cup v$, the product of u and v , as follows. (We assume that $s+t \leq 2r+1$, since otherwise $u \cup v = 0$.)

Case 1. If $s+t \leq r$, then $u \in A_{sp} = \wedge^s V$, $v \in A_{tp} = \wedge^t V$, and we define $u \cup v \in A_{(s+t)p} = \wedge^{s+t} V$ to be $u \wedge v$.

Case 2. If $s \leq r, t \leq r$ and $s+t \geq r+1$, then we define $u \cup v \in A_{(s+t)p} = (\wedge^{2r+1-s-t} V)^*$ by the equation $(u \cup v)(w) = F(u \wedge v \wedge w)$ for $w \in \wedge^{2r+1-s-t} V$.

Case 3. If $s \leq r, t \geq r+1$ and $s+t \leq 2r+1$, then $u \in A_{sp} = \wedge^s V$, $v \in A_{tp} = (\wedge^{2r+1-t} V)^*$, and we define $u \cup v \in A_{(s+t)p} = (\wedge^{2r+1-s-t} V)^*$ by the equation $(u \cup v)(w) = v(w \wedge u)$ for $w \in \wedge^{2r+1-s-t} V$.

Case 4. If $s \geq r+1, t \leq r$ and $s+t \leq 2r+1$, then $u \in A_{sp} = (\wedge^{2r+1-s} V)^*$, $v \in A_{tp} = \wedge^t V$, and we define $u \cup v \in A_{(s+t)p} = (\wedge^{2r+1-s-t} V)^*$ by the equation $(u \cup v)(w) = u(v \wedge w)$ for $w \in \wedge^{2r+1-s-t} V$.

Routine calculation shows that this product makes A a commutative graded algebra satisfying Poincaré duality and conditions (i) except that $A_{pq} = \mathbb{Q}^*$. By the canonical isomorphism $f : \mathbb{Q}^* \rightarrow \mathbb{Q}$ (i.e. for $w \in \mathbb{Q}^*$, $f(w) = w(1)$), we can identify A_{pq} with $\mathbb{Q}$, then it is easy to see that condition (ii) holds.

If p is even, we can just change all $\wedge$ in the above construction by $\vee$. $\square$

From [Su], Theorem (13.2), we have

Theorem 4.3. *Any commutative graded algebra A over $\mathbb{Q}$ satisfying Poincaré duality and $A_1 = 0$ may be realized as the rational cohomology ring of a simply-connected, closed smooth manifold of dimension n with possibly one singular point when $n = 4k$. Moreover, when $n = 4k$, if $A_{2k} = 0$, then we can remove the singularity in a realizing manifold.*

Proof of Theorem 4.1*. Suppose such an $(n+1)$ -manifold W exists. Since n is a composite number and is not a power of 2, we have a factorization $n = pq$, where p is an integer ≥ 2 and q is an odd integer ≥ 3 . By Proposition 3.1, we can select an integer $m \geq 3\beta_{p+1}(W)$ such that X_m can be constructed in $(\wedge^q \mathbb{R}^m)^*$ (when p is odd) or $(\vee^q \mathbb{R}^m)^*$ (when p is even). Since X_m is proper closed, we can pick an $F \notin X_m$ such that $F(e_{i_1}, \dots, e_{i_q}) \in \mathbb{Q}$ for any $1 \leq i_1, \dots, i_q \leq m$.

Taking $V = \mathbb{Q}^m$, by Proposition 4.2, there exists a commutative graded algebra A satisfying Poincaré duality and conditions (i) and (ii). $p \geq 2$ guarantees that $A_1 = 0$. By Theorem 4.3, there exists a simply-connected, closed, smooth n -manifold M such that the rational cohomology ring $H^*(M; \mathbb{Q})$ is isomorphic to A . (When $n = 4k$, $A_{2k} = 0$ as $2k$ is not an integral multiple of p . Thus we can remove the singularity in a realizing manifold.) Denote this isomorphism by $h_{\mathbb{Q}}$, and define $h_{\mathbb{R}}$ to be

$$h_{\mathbb{Q}} \otimes \text{id} : H^*(M; \mathbb{R}) = H^*(M; \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{R} \rightarrow A \otimes_{\mathbb{Q}} \mathbb{R}.$$

Clearly $h_{\mathbb{R}}$ gives isomorphisms $H^p(M; \mathbb{R}) \rightarrow \mathbb{R}^m$ and $H^n(M; \mathbb{R}) \rightarrow \mathbb{R}$. Condition (ii) in Proposition 4.2 can be transformed to

$$F(h_{\mathbb{R}}(x_1), \dots, h_{\mathbb{R}}(x_q)) = h_{\mathbb{R}}(x_1 \cup \dots \cup x_q). \quad (5)$$

Note that $\beta_p(M) = m \geq 3\beta_{p+1}(W)$, hence $\frac{1}{2}(\beta_p(M) - \beta_{p+1}(W)) \geq \frac{m}{3}$. Since M can be topologically flat embedded into W , by Proposition 2.1, there exists a subspace V of $H^p(M; \mathbb{R})$ and a linear transformation $\varphi : V \rightarrow H^p(M; \mathbb{R})$ with no fixed non-zero vectors such that

$$\dim V \geq \frac{1}{2}(\beta_p(M) - \beta_{p+1}(W)) \geq \frac{m}{3}$$

and for any $x_1, \dots, x_q \in V$,

$$x_1 \cup \dots \cup x_q = \varphi(x_1) \cup \dots \cup \varphi(x_q). \quad (6)$$

Let $U = h_{\mathbb{R}}(V)$. Define $\psi : U \rightarrow \mathbb{R}^m$ by $\psi(y) = (h_{\mathbb{R}} \circ \varphi \circ h_{\mathbb{R}}^{-1})(y)$ for $y \in U$. Then $\dim U \geq \frac{m}{3}$ and $\psi : U \rightarrow \mathbb{R}^m$ is a linear map with no fixed non-zero vectors. By equations (5) and (6), for any $y_1, \dots, y_q \in U$, we have

$$F(y_1, \dots, y_q) = F(\psi(y_1), \dots, \psi(y_q)).$$

Thus F is special. But $F \notin X_m$, a contradiction. $\square$

5 Embedding of 4-manifolds into 5-manifolds

We first list theorems about the classification of simply-connected 4-manifolds according to their intersection forms. More details can be found in [Ki] or [GS].

For any compact, connected, oriented 4-manifold M , the cup product

$$\cup : H^2(M, \partial M) \times H^2(M, \partial M) \rightarrow H^4(M, \partial M), (\alpha, \beta) \mapsto \alpha \cup \beta$$

gives a symmetric bilinear form Q_M over $\mathbb{Z}$ on $H_2(M)$ through the isomorphisms $H^2(M, \partial M) \cong H_2(M)$ and $H^4(M, \partial M) \cong \mathbb{Z}$. Clearly, $Q_M(a, b) = 0$ if a or b is a torsion element. So Q_M descends to a symmetric bilinear form on $H_2(M)/\text{Torsion} \cong \mathbb{Z}^r$.

For a given integral symmetric bilinear form Q on a finitely generated free abelian group A , the *rank*, *signature* and *parity* of Q are defined as follows: the rank $\text{rk}(Q)$ of Q is the dimension of A . Extend and diagonalize Q over $A \otimes_{\mathbb{Z}} \mathbb{R}$, the number of positive entries and the number of negative entries are denoted by b_2^+ and b_2^- respectively, and the difference $b_2^+ - b_2^-$ is the signature $\sigma(Q)$ of Q . Finally, Q is even if $Q(a, a)$ is even for any $a \in A$ and Q is odd otherwise.

By choosing a basis for A , one can express Q as a symmetric matrix. The determinant of this matrix does not depend on the choice of the basis, and thus we denote it by $\det Q$. If $\det Q = \pm 1$, we say Q is *unimodular*. It follows from Poincaré duality that Q_M is unimodular if M is closed. We say Q is *indefinite* if both b_2^+ and b_2^- are positive.

Theorem 5.1. (cf. [GS], Theorem 1.2.21) *Suppose Q is an integral, indefinite, unimodular symmetric bilinear form. If Q is odd, then it is isomorphic to $b_2^+ \langle +1 \rangle \oplus b_2^- \langle -1 \rangle$; if Q is even, then it is isomorphic to $\frac{\sigma(Q)}{8} E_8 \oplus \frac{\text{rk}(Q) - |\sigma(Q)|}{2} H$, where E_8 is an even form with $\text{rk}(E_8) = \sigma(E_8) = 8$ and $H = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$.*

This theorem classifies indefinite unimodular forms, while the next theorem is a classification result for simply-connected closed 4-manifolds.

Theorem 5.2. (cf. [Fr]) *For every integral, unimodular symmetric bilinear form Q , there exists a simply-connected, closed, topological 4-manifold M such that $Q_M \cong Q$.*

If Q is even, this manifold is unique up to homeomorphism. If Q is odd, there are exactly two different homeomorphism types of such manifolds. At most one of these two homeomorphism types carries a smooth structure. Consequently, simply-connected, closed, smooth 4-manifolds are determined up to homeomorphism by their intersection forms.

In the odd case, the two manifolds are distinguished by their Kirby-Siebenmann triangulation invariants $\text{KS} \in \mathbb{Z}_2$. One has non-zero invariant and cannot be smooth, while the other has zero invariant and is stably smoothable.

For a compact, connected, topological 4-manifold M , its Kirby-Siebenmann invariant (cf. [KS]) $\text{KS}(M) \in \mathbb{Z}_2$ is the stable smoothing obstruction. This invariant is additive in the following sense (cf. [FQ], Section 10.2B): If there is a homeomorphism of boundaries $\partial M \cong \partial N$, then $\text{KS}(M \cup_{\partial} N) = \text{KS}(M) + \text{KS}(N)$. Combined with the fact that D^4 is smoothable, we have

Theorem 5.3. *Suppose M_1 and M_2 are two connected, closed, topological 4-manifolds, then*

$$\text{KS}(M_1 \# M_2) = \text{KS}(M_1) + \text{KS}(M_2).$$

Next, we give a construction that will be used many times later.

Lemma 5.4. *For any oriented, closed, connected 4-manifold M , there exists an oriented, closed, connected 5-manifold W such that for any positive integer r , $\#rM$ can be topologically flat embedded into W .*

Proof. We first construct the double connected sum of M . Let D_1^4 and D_2^4 be two disjoint embedded 4-balls in M , and let X be

$$(M - \text{int}(D_1^4) - \text{int}(D_2^4)) \cup_{\varphi} (M - \text{int}(D_1^4) - \text{int}(D_2^4)),$$

where φ is an orientation-reversing homeomorphism of $\partial D_1^4 \amalg \partial D_2^4$.




Figure 1

Clearly, X consists of 2 copies of $M - 2 \text{int}(D^4)$ and 2 copies of $S^3 \times I$. We denote these two kinds of pieces by dots and arcs respectively as shown in Figure 1.

Next, we apply surgeries on $X \times S^1$ by cutting out two disjoint copies of $(S^3 \times I) \times I = S^3 \times D^2$ and replacing each of them by $D^4 \times S^1$. The new manifold is denoted by W .




Figure 2

In Figure 2, W is depicted as a torus, where the X and the S^1 factors correspond to meridian and longitude respectively. In this picture, each point in the two thickened longitudes represents $M - 2 \text{ int}(D^4)$, any other point outside the squares represents S^3 , and the two shaded squares are the surgery region.

We claim that for any positive integer r , $\#rM$ can be embedded into W .




Figure 3

In Figure 3, a long curve is drawn with endpoints in the surgery region and r dots on it. (The left picture is for even r , while the right picture is for odd r .) Each dot represents $M - 2 \text{ int}(D^4)$ and each arc in the curve that does not meet the surgery region is $S^3 \times I$. For an arc that goes into the surgery region, its intersection with the boundary of the surgery region is S^3 , which bounds a D^4 in W . Hence the first and the last arc in the curve actually both denote D^4 . Now this curve gives an embedding of $\#rM$ in W . $\square$

Now we can state and prove our results.

Theorem 5.5. (a) *There exists an oriented, connected, closed 5-manifold W , such that for any simply-connected, closed 4-manifold M with indefinite intersection form, M can be topologically flat embedded into W .*

(b) *Any simply-connected, compact topological 4-manifold M with $\partial M \neq \emptyset$ admits a topologically flat embedding into $S^2 \tilde{\times} S^3$.*

Proof. (a) Clearly, $\mathbb{CP}^2$ and $\overline{\mathbb{CP}}^2$ are smooth manifolds with intersection form $\langle +1 \rangle$ and $\langle -1 \rangle$ respectively. And there are non-smoothable manifolds X_1 and X_{-1} that also have intersection form $\langle +1 \rangle$ and $\langle -1 \rangle$ respectively. Then

$$X_1 \# (p-1)\mathbb{CP}^2 \# q\overline{\mathbb{CP}}^2 \text{ and } X_2 \# p\mathbb{CP}^2 \# (q-1)\overline{\mathbb{CP}}^2$$

have intersection form $p\langle +1 \rangle \oplus q\langle -1 \rangle$ and non-zero KS invariant by Theorem 5.3 if the expressions are meaningful. Now it follows from Theorems 5.1 and 5.2 that any simply-connected, oriented, closed 4-manifold M with odd, indefinite intersection form is homeomorphic to

$$k_1 X_1 \# k_2 X_{-1} \# k_3 \mathbb{CP}^2 \# k_4 \overline{\mathbb{CP}}^2,$$

for certain nonnegative integers k_i .

Let X_{E_8} and X_{-E_8} be 4-manifolds with intersection form E_8 and $-E_8$, respectively. Clearly $S^2 \times S^2$ has intersection form H . Then it still follows from Theorems 5.1 and 5.2 that any simply-connected, oriented, closed 4-manifold M with even, indefinite intersection form is homeomorphic to

$$l_1 X_{E_8} \# l_2 X_{-E_8} \# l_3 (S^2 \times S^2)$$

for certain nonnegative integers l_i .

Choose M to be $X_1, X_{-1}, \mathbb{CP}^2, \overline{\mathbb{CP}}^2, X_{E_8}, X_{-E_8}$ and $S^2 \times S^2$ respectively and construct corresponding 5-manifolds $W_1, W_2, \dots, W_7$ by Lemma 5.4. It follows from the claim below that the desired manifold W can be chosen as $W_1 \# W_2 \# \dots \# W_7$.

Claim. If there are topologically flat embeddings $M_1 \hookrightarrow W_1$ and $M_2 \hookrightarrow W_2$, then $M_1 \# M_2$ can be topologically flat embedded into $W_1 \# W_2$.

Proof of the claim. Choose D_i^4 in M_i . Since $M_i \times I$ is a part of W_i , $D_i^4 \times I$ is a 5-ball in W_i . Cut out these two 5-balls from W_i respectively and glue their boundaries, then we have an embedding of $(M_1 \# M_2) \times I$ into $W_1 \# W_2$.

(b) We glue M and $-M$ together by the identity map between ∂M and $\partial(-M)$. The resulting manifold is usually called the double of M and denoted by DM . Clearly DM is a simply-connected, closed 4-manifold with $\sigma(DM) = \sigma(M) + \sigma(-M) = 0$ (cf. [Ki], Chapter II. Theorem 5.3) and $\text{KS}(DM) =$

$\text{KS}(M) + \text{KS}(-M) = 0$. It follows from Theorem 5.1 that the intersection form Q of DM is either

$$k\langle +1 \rangle \oplus k\langle -1 \rangle \text{ or } kH,$$

depending on whether Q is odd or even.

Hence the intersection form of DM is isomorphic to that of either $\#k\mathbb{CP}^2 \# k\overline{\mathbb{CP}}^2$ or $\#kS^2 \times S^2$. It is well known that

$$\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2 \cong S^2 \tilde{\times} S^2, \quad S^2 \tilde{\times} S^2 \# S^2 \tilde{\times} S^2 \cong S^2 \times S^2 \# S^2 \tilde{\times} S^2.$$

Thus by $\text{KS}(DM) = 0$ and Theorem 5.2, DM is homeomorphic to either $\#kS^2 \times S^2$ or $\#(k-1)S^2 \times S^2 \# S^2 \tilde{\times} S^2$.

$S^2 \tilde{\times} S^2$ can be embedded into $S^2 \tilde{\times} S^3$ because the inclusion $\text{SO}(3) \hookrightarrow \text{SO}(4)$ induces isomorphism $\pi_1(\text{SO}(3)) \cong \pi_1(\text{SO}(4)) \cong \mathbb{Z}_2$. And $S^2 \times S^2$ can be embedded into $\mathbb{R}^5$ through the standard inclusion $S^2 \times D^3 \hookrightarrow \mathbb{R}^5$. Therefore by the claim in the proof of (a), both $\#kS^2 \times S^2$ and $\#(k-1)S^2 \times S^2 \# S^2 \tilde{\times} S^2$ can be embedded into $S^2 \tilde{\times} S^3$. $\square$

Acknowledgements. The authors would like to thank Jianzhong Pan for informing us the reference of Sullivan's work [Su2] and for other interesting talks. The first two authors are partially supported by grant no. 10201003 of NSFC and a grant of MSTC.

References

- [Di] **F. Ding**, *Smooth structures on some open 4-manifolds*, Topology 36 (1997), no. 1, 203–207.
- [EL] **A. L. Edmonds, C. Livingston**, *Embedding punctured lens spaces in four-manifolds*, Comment. Math. Helv. 71 (1996), no. 2, 169–191.
- [Fa1] **F. Fang**, *Embedding 3-manifolds and smooth structures of 4-manifolds*, Topology Appl. 76 (1997), no. 3, 249–259.
- [Fa2] **F. Fang**, *Smooth structures on $\Sigma \times \mathbf{R}$* , Topology Appl. 99 (1999), no. 1, 123–131.
- [Fr] **M. H. Freedman**, *The topology of four-dimensional manifolds*, J. Differential Geom. 17 (1982), no. 3, 357–453.
- [FQ] **M. H. Freedman, F. Quinn**, *Topology of 4-manifolds*, Princeton University Press, Princeton, NJ, 1990.
- [GS] **R. E. Gompf, A. I. Stipsicz**, *4-manifolds and Kirby calculus*, Graduate Studies in Math. 20, Amer. Math. Soc., Providence, RI, 1999.
- [Hi] **M. W. Hirsch**, *The imbedding of bounding manifolds in euclidean space*, Ann. of Math. (2) 74 (1961), 494–497.

- [HJ] **R. A. Horn, C. R. Johnson**, *Matrix analysis*, Cambridge University Press, Cambridge, 1990.
- [Ka] **A. Kawauchi**, *The imbedding problem of 3-manifolds into 4-manifolds*, Osaka J. Math. 25 (1988), no. 1, 171–183.
- [Ki] **R. C. Kirby**, *The topology of 4-manifolds*, Lecture Notes in Math. 1374, Springer-Verlag, Berlin, 1989.
- [KS] **R. C. Kirby, L. C. Siebenmann**, *Foundational essays on triangulations and smoothings of topological manifolds*, , Ann. of Math. Studies 88, Princeton University Press, Princeton, NJ, 1977.
- [La] **S. Lang**, *Algebra*, Graduate Texts in Math. 211, Springer-Verlag, New York, 2002.
- [Ro] **V. A. Rohlin**, *The embedding of non-orientable three-manifolds into five-dimensional Euclidean space*, Soviet Math. Dokl. 6 (1965), 153–156.
- [Sh] **R. Shaw**, *Linear algebra and group representations, Vol. II*, Academic Press, London-New York, 1983.
- [Shi] **T. Shiomi**, *On imbedding 3-manifolds into 4-manifolds*, Osaka J. Math. 28 (1991), no. 3, 649–661.
- [Sp] **E. H. Spanier**, *Algebraic topology*, Springer-Verlag, New York, 1966.
- [Su] **D. Sullivan**, *Infinitesimal computations in topology*, Inst. Hautes Études Sci. Publ. Math. 47 (1977), 269–331 (1978).
- [Wa] **C. T. C. Wall**, *All 3-manifolds imbed in 5-space*, Bull. Amer. Math. Soc. 71 (1965), 564–567.
- [WZ] **S. Wang, Q. Zhou**, *How to embed 3-manifolds into 5-space*, Adv. in Math. (China) 24 (1995), no. 4, 309–312.

LMAM AND SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, P.R.CHINA

E-mail address: dingfan@math.pku.edu.cn

LMAM AND SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, P.R.CHINA

E-mail address: wangsc@math.pku.edu.cn

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT BERKELEY, CA 94720, USA

E-mail address: jgyao@math.berkeley.edu